

If you are thinking about having chorionic villus sampling or amniocentesis, talk it over with your doctor. They can make a referral to the MFM clinic if you want more information.

通常在妊娠第二期（即妊娠第16至20週期間）進行，方法是利用一根細針抽取少量胎盤中的羊水，以檢測胎兒有否脊柱裂（一種神經管閉合缺漏的畸形）或異常的染色體組型。孕婦可能基於多種不同原因而需要接受此項檢測，包括下列情況：

- 年滿35歲或以上的孕婦；
- 有家族遺傳病史；
- 唐氏綜合症早期篩查顯示懷有高危胎兒；
- 超聲波測試中發現問題。

抽胎水導致流產的機會率約0.5%。

如覺得需要接受絨毛活檢或抽胎水檢查，宜跟醫生詳細討論。如欲進一步了解有關這兩項檢查，可由醫生轉介至母胎醫學診所。

What do I do if I have a Down's Syndrome fetus? 如果我的胎兒有唐氏綜合症該怎麼辦？

Your specialist will discuss with you the likely outcome and associated difficulties, and can provide information about support services in Hong Kong.

You can discuss with your obstetrician the alternative of termination of pregnancy, which in Hong Kong is considered both legal and safe if performed not later than 24 weeks gestation.

萬一不幸確診胎兒有唐氏綜合症，醫生將會與你詳細討論和分析，包括有可能面對的困難和處理方法，並提供有關服務的資料。

在香港，妊娠24週內終止妊娠是合法的，有關手術亦是安全的，請跟婦產科醫生討論有關細節。

Appointment & Enquiries 預約及查詢

Please telephone our Registration Office on

請致電本院登記處預約時間及計劃

Tel 電話：(852) 2276 7022




Like us on

Hong Kong Adventist Hospital - Tsuen Wan



Map 路線圖



Bus 巴士

39A Tsuen Wan West Station
39M Tsuen Wan MTR
30X Whampoa Gardens
30 Cheung Sha Wan
A38 Hong Kong International Airport

To the Hospital Via Castle Peak Road:
+ Alight at Tsuen King Circuit Flyover
+ Take the bus in front of Tsuen King Circuit Police Station

荃灣西站
荃灣港鐵站
黃埔花園
長沙灣
香港國際機場

所有行經青山公路荃灣段之巴士：
+ 於荃景圍天橋站下車，
+ 然後步行至荃景圍警署轉乘巴士到醫院。

Minibus 專線小巴

95M Tsuen Wan MTR
95 Nina Tower
84 Tsuen Wan Chung On Street

荃灣港鐵站
如心廣場
荃灣眾安街

24小時急診中心 Hour Urgent Care Center

Hong Kong Adventist Hospital - Tsuen Wan
香港港安醫院·荃灣

Address 地址：199 Tsuen King Circuit, Tsuen Wan, N.T.
新界荃灣荃景圍199號

Telephone 電話：(852) 2275 6688

Fax 傳真：(852) 2275 6767

Website 網站：www.twah.org.hk



Hong Kong Adventist Hospital - Tsuen Wan reserves the right to update the fee schedule, brochure, terms and conditions. Any change of fee schedule will be announced and notices will be published in advance, in accordance with the statutory notice period. Any other changes except the fee schedule may be made at any time as it sees fit without prior notice, you are advised to check the latest information before using our services. For the latest information, please visit our website at www.twah.org.hk
香港港安醫院—荃灣保留修訂收費表、單張內容、條款及細則之權利。任何的收費表調整將會根據法定的通知期提前發出通告和作出宣布。其他與收費表以外的相關修訂，則可能不作另行通知，閣下於使用本院服務前查閱最新資訊。
如欲查閱最新資料，請瀏覽本院網站www.twah.org.hk。

Valid date 有效期至：2025/12/31

N-2504

Adventist 港
Health 安

Hong Kong Adventist Hospital - Tsuen Wan

香港港安醫院·荃灣



Maternal Fetal Medicine Clinic 母胎醫學診所



Extending the Healing Ministry of Christ
延續基督的醫治大能



The Maternal Fetal Medicine Clinic provides various antenatal screening and diagnostic tests. We hope our service can add more information for the preparation of the birth for your precious baby.

- Antenatal Down's Syndrome Screening Test
- Chorionic Villus Sampling
- Amniocentesis
- Second Trimester Ultrasound Scanning
- Pre-pregnancy Counseling

母胎醫學診所提供多種產前掃描篩查和診斷檢測，誠意為大家提供更多資料，以為迎接新生命來臨作最佳準備。

- 產前唐氏綜合症普查
- 絨毛活檢
- 抽胎水
- 孕中期詳細超聲波掃描
- 產前輔導

What is Down's Syndrome? 甚麼是唐氏綜合症?

Down's Syndrome is a condition resulting from a genetic abnormality, in which an extra chromosome is present. (There are three no.21 chromosomes instead of the usual 2, giving a total of 47 chromosomes instead of the normal 46) The distribution of chromosomes may go wrong around the time of conception. Baby may receive an extra chromosome from either parent, resulting in an abnormal number of chromosomes.

Apart from their characteristic appearance, the affected individuals may have a degree of mental handicap. Associated abnormalities may also be present, of which heart defects are the most common.

Down's syndrome occurs must frequently by chance and is seldom caused by parental inheritance.

The risks of having a Down's Syndrome baby increases with maternal age. Overall, the risk is 1:700, but for women of 35 years, the risk is 1:400, and for women of 40 years or more, the risk is 1:100.

With multiple pregnancies, the risk may differ, and the mother should consult with her obstetrician.



唐氏綜合症源於胎兒的染色體異常，亦即他第二十一組染色體較正常人多了一條，共有三條。正常胎兒的細胞中含有二十三組染色體，每組兩條，分別來自父親和母親。但在受孕期間，染色體的分佈可能發生錯誤。胎兒有可能從雙親中任何一方得到一條額外的染色體，致染色體數目異常。染色體中包含了控制人體成長和機能的基因，故當分佈出錯，就會影響胎兒發育。這影響可以是多方面的，包括外貌、智力、身體結構等。

唐氏綜合症兒童除了出現外貌特徵外，往往有不同程度的學習障礙，並可能會有先天性心臟病等問題。

大多數唐氏綜合症屬偶發性，絕少是受父母遺傳，機會率為七百分之一。雖然任何年齡的孕婦都會有機會懷上唐氏綜合症胎兒，但是機會率隨孕婦年齡增加而升高。例如，一名35歲孕婦懷唐氏綜合症胎兒的機會是四百分之一，而40歲孕婦則為百分之一。

如屬多胎妊娠，胎兒患唐氏綜合症的機會率有可能會不同，故宜跟婦產科醫生查詢。

Prenatal Screening for Down's Syndrome? 甚麼是「產前唐氏綜合症普查」?

The mother's age is significant. Ultrasound is used to measure the nuchal fold of the fetus between the 11th - 13th week of pregnancy. The inclusion of quantitative analysis of HCG and PAPP-A in maternal blood will increase the detection rate to 85%. Results of these screening tests are available within 24 hours.

「產前唐氏綜合症普查」是根據母親年齡推測懷唐氏綜合症胎兒的機會率，再結合超聲波標記和母親血清學測試進行風險估計。胎兒的超聲波檢查是測量頸皮厚度，須在妊娠第11至第13週期間進行，如果將母親血液中的標記物HCG和PAPP-A都納入分析，檢出率可達85%。有關結果可在接受測試後二十四小時內知道。



Interpretation of Screening Test Results? 如何解讀測試結果?

The test results are reported as either 'positive = high risk' or 'negative = low risk'. Patients assessed as 'high risk' are offered amniocentesis or chorionic villi sampling for definitive diagnosis.

Note: The majority of patients who are 'positive' on screening, do not in fact have a Down's Syndrome foetus. A screening test 'negative' does not entirely rule out the possibility of Down's Syndrome, but with this result, the risk is judged very low.

測試結果可分為陰性（低危）和陽性（高危）兩種，如結果為陽性，即胎兒患上唐氏綜合症的風險較高，須接受診斷測試，即抽胎水或絨毛活檢。如結果為陰性，即胎兒患上唐氏綜合症的風險較低。

備註：大部份測試結果為「陽性」的孕婦，實際上並未懷有唐氏綜合症胎兒。另一方面，即使結果為「陰性」，並不代表已確診胎兒100%不會患上唐氏綜合症。

What is Chorionic Villi Sampling? 甚麼是絨毛活檢?

This procedure is performed in first trimester (11-14 weeks of pregnancy). A needle is used to take the placental tissue. The sample is tested for fetal karyotype. Chorionic villus sampling can cause a miscarriage rate in about one out of 100 women.

絨毛活檢通常在妊娠第一期（即妊娠第11至14週期間）進行，方法是利用一根細針抽取少量胎盤組織，以檢測胎兒的染色體組型。接受此檢驗手術而導致流產的機會率為1%。

What is Amniocentesis? 甚麼是抽胎水?

This procedure is performed in second trimester (16-20 weeks of pregnancy). A needle is used to take a small sample of the amniotic fluid that surrounds the fetus. The fluid is tested to see if the fetus had spina bifida (a type of open neural tube defect) or abnormal fetal karyotype (including Down's Syndrome). Women have amniocentesis for many different reasons. You may wish to have an amniocentesis if:

- You will be 35 years or more when your baby is due
- You have an inherited disease present in your family
- Your antenatal Down screen has put you at an increased risk
- A problem has been identified on your ultrasound

Amniocentesis can cause a miscarriage in about one out of 200 women.



Maternal Fetal Medicine Clinic

Maternal Fetal Medicine Clinic



Examination Items

Fee

Antenatal Down's Syndrome Screening Tests

<input type="checkbox"/> SAFET21	\$4500
<input type="checkbox"/> SAFET21 Advanced Profile	\$6000
<input type="checkbox"/> First Trimester Fetal Nuchal Translucency	\$1700

Detailed Ultrasound Scan

<input type="checkbox"/> Second Trimester Detailed Ultrasound Scan	
● Single	\$3750
● Twin	\$7500
<input type="checkbox"/> Fetal Growth Assessment + Doppler Study	\$2370
<input type="checkbox"/> Third Trimester Detail Ultrasound Scan	
● Single	\$2000
● Twin	\$4000

Prenatal Diagnostic Tests

<input type="checkbox"/> Amniocentesis	\$4500
<input type="checkbox"/> Chorionic Villus Sampling	\$4500

Optional Items

Genetic Test

● Chromosome Microarray	\$4510
● Karyotype	\$2970
● PCR	\$1650

<input type="checkbox"/> Prenatal Counseling	\$2150
---	--------

<input type="checkbox"/> Prepregnancy Counseling	\$2150
---	--------

+ Important information

+ The price list is applicable for singleton pregnancy.

+ Appointment & Enquiries

For further information or appointment with our Maternal Fetal Medicine Clinic, please contact us on the telephone number below, or visit our website.

Tel : (852) 2276 7022

Website : www.twah.org.hk

Valid date : 2025/12/31

N-2504



Maternal Fetal Medicine

檢查計劃

費用

產前唐氏綜合症普查

■ 敏兒安	\$4500
■ 進階版敏兒安	\$6000
■ 孕早期胎兒後頸皮厚度量度	\$1700

詳細超聲波掃描

■ 孕中期詳細超聲波掃描	
● 單胎	\$3750
● 雙胞胎	\$7500
■ 胎兒生長評估 + 多普勒超聲波胎兒血管血流量度	\$2370
■ 懷孕後期詳細超聲波掃描	
● 單胎	\$2000
● 雙胞胎	\$4000

產前診斷測試

■ 抽胎水	\$4500
■ 絨毛活檢	\$4500

另加項目

遺傳學

● 染色體微陣列分析	\$4510
● 染色體核型分析	\$2970
● 聚合酶鏈反應	\$1650

■ 產前諮詢	\$2150
■ 懷孕前諮詢	\$2150

注意事項

+ 除特別註明外，上述價錢以單胎妊娠計算。

預約及查詢

如欲進一步了解，或預約本院母胎醫學診所，請與我們聯絡，或瀏覽下列網站。

電話：(852) 2276 7022

網址：www.twah.org.hk

